

# **Identification and Antioxidant Activity Study of Bioactive Compounds in Areca Nut Oil**

Mingjun Zhu

Hainan Tropical Ocean College, Sanya, 572022, China.

**Abstract:** This study aims to identify the bioactive compounds in areca nut oil and evaluate their antioxidant properties. Utilizing advanced chromatography-mass spectrometry techniques and bioactivity detection methods, a variety of compounds with potential health benefits were successfully isolated and identified from areca nut oil. The results indicate that these compounds possess significant antioxidant capabilities, effectively scavenging free radicals and reducing oxidative stress, offering new insights for the development of novel natural antioxidants and the prevention of chronic diseases.

**Keywords:** Areca Nut Oil; Bioactive Compounds; Antioxidant; Chromatography-Mass Spectrometry; Free Radicals

**Introduction:** Areca nut oil, extracted from areca nut seeds, has historically been used in traditional medicine and cooking. In recent years, as the role of natural products in health care and disease prevention has become increasingly prominent, areca nut oil and its bioactive components have attracted widespread scientific attention. Antioxidants play a crucial role in preventing cell damage and chronic diseases caused by free radicals. However, the specific bioactive

compounds present in areca nut oil and their potential health benefits require further research and exploration.

## **1Chemical Component Analysis of Areca Nut Oil**

### **1.1 Extraction and Separation Techniques**

Areca nut oil, extracted from the mature seeds of the areca nut, has attracted widespread attention in the food and pharmaceutical industries due to its variety of bioactive components. Traditional methods of extracting areca nut oil, such as cold pressing and hot pressing, may not effectively preserve the integrity and stability of its bioactive components. Therefore, this study employed supercritical carbon dioxide extraction techniques to maximize the retention of active components in areca nut oil.

Initially, collected mature areca nut seeds were cleaned and the outer hard shell removed. The seeds were then cut into small pieces using a cutting machine to increase the surface area during the extraction process and reduce the amount of solvent needed. The chopped seeds were placed in a supercritical fluid extractor, and supercritical carbon dioxide under preset conditions was introduced. In its supercritical state, carbon dioxide has low viscosity and high diffusion rates, enabling it to penetrate the porous structure of plant cells and effectively dissolve fats and other lipophilic components.<sup>[1]</sup>

Throughout the extraction process, temperature and pressure

conditions were strictly controlled to ensure optimal extraction efficiency without damaging sensitive bioactive molecules. The extraction continued for several hours until a sufficient quantity of oil was extracted from the raw material. After extraction, the supercritical carbon dioxide was returned to a gaseous state by reducing pressure and temperature, thus separating out the areca nut oil.

Upon completion of this stage, a batch of dark brown areca nut oil was obtained. To further purify the oil, removing possible impurities and colorants, methods such as molecular distillation and activated carbon adsorption were used. Molecular distillation occurs at lower temperatures, reducing the likelihood of thermal decomposition, while activated carbon adsorption effectively removes residual impurities and unwanted colorants. After purification, the areca nut oil appeared clearer, turning a golden-yellow color, laying a good foundation for subsequent analysis and assessment work. [2]

## **1.2 Identification Using Chromatography-Mass Spectrometry**

To delve deeper into the chemical composition of areca nut oil, especially its bioactive components, this study employed liquid chromatography-mass spectrometry (LC-MS) and gas chromatography-mass spectrometry (GC-MS) techniques for detailed analysis.

Firstly, the purified areca nut oil samples were suitably diluted and

pretreated for chromatographic analysis. The samples were then separated using specially selected chromatographic columns based on the types of compounds to be separated, ensuring efficient and accurate separation.[3] In LC-MS analysis, particular attention was paid to polyphenols and other highly polar compounds, while GC-MS was primarily used to analyze fatty acids and volatile components.

During the chromatographic separation process, compounds were separated based on their interactions between the mobile and stationary phases. Once separated, these compounds entered the mass spectrometer for further identification and quantification. Mass spectrometry analysis involved measuring the mass-to-charge ratio ( $m/z$ ) of ionized compounds, crucial for determining the molecular structure of unknown compounds.

Through comprehensive analysis of LC-MS and GC-MS results, we successfully identified a range of key components in areca nut oil, including several polyunsaturated fatty acids, liposoluble vitamins, carotenoids, phenolic compounds, and other bioactive small molecules. Notably, some rare polyphenolic compounds, uncommon in plant oils but exhibiting strong antioxidant and anti-inflammatory properties, were of particular interest.[4]

In conclusion, through the application of supercritical carbon dioxide extraction and chromatography-mass spectrometry techniques,

this study not only successfully extracted and identified the main bioactive components in areca nut oil but also provided a solid foundation for further biological evaluation and potential health applications of these components. In future work, these components will undergo more in-depth functional studies and activity validations, potentially playing a larger role in the fields of food science and pharmaceutical research. [5]

## **2Study of Antioxidant Properties of Bioactive Compounds in Areca Nut Oil**

### **2.1 Methods of Assessing Antioxidant Activity**

The presence of various bioactive compounds in areca nut oil indicates its potential health benefits, particularly its antioxidant properties. Oxidative stress is a key factor leading to cellular damage, aging, and various diseases. This chapter aims to explore the ability of specific active components in areca nut oil to combat free radicals and inhibit oxidative stress through a series of experimental methods.

The experiments first employed a variety of established and widely accepted in vitro antioxidant assessment methods. These assessment methods include DPPH free radical scavenging assay, ABTS scavenging assay, reducing power assay, iron chelating activity test, and lipid peroxidation inhibition assay.

DPPH Free Radical Scavenging Assay: The DPPH

(1,1-Diphenyl-2-picrylhydrazyl) free radical scavenging assay is a common method used to assess the antioxidant's capacity to scavenge the stable free radical DPPH. In this assay, DPPH free radicals react with potential antioxidants, leading to a color change in the DPPH solution from purple to yellow. This color change is quantified by measuring the reduction in absorbance at 517 nm. The antioxidant capacity of the sample is evaluated by measuring the half maximal inhibitory concentration (IC<sub>50</sub>), i.e., the concentration of the extract or compound needed to scavenge 50% of DPPH free radicals. In our study, the DPPH assay provided preliminary information on the free radical scavenging ability of antioxidant compounds in areca nut oil. [6]

**ABTS Scavenging Assay:** The ABTS scavenging assay uses the ABTS free radical system (2,2'-Azino-bis[3-ethylbenzothiazoline-6-sulfonic acid]-6-diethylbenzothiazole). The principle is similar to the DPPH assay but uses ABTS free radicals. These radicals react with antioxidants, leading to a change in solution color, which is quantified by measuring the reduction in absorbance at 734 nm. The ABTS assay is an effective method to assess the activity of both water-soluble and lipid-soluble antioxidants, as ABTS free radicals are soluble in both water and organic solvents. In our areca nut oil samples, this method helped us assess the scavenging ability of different compounds against ABTS radicals.

**Reducing Power Assay:** The reducing power assay is based on the

electron donation ability of the sample, i.e., the reducing capacity of the antioxidant. In this method, antioxidants demonstrate their reducing power by converting  $\text{Fe}^{3+}$  to  $\text{Fe}^{2+}$ . The extent of  $\text{Fe}^{2+}$  formation is measured by its complexation reaction with reducing agents like potassium ferricyanide and measured by the absorbance of the formed complex at 700 nm. In our study, this assessment method allowed us to determine which components in areca nut oil have higher reducing capacities, an important component of their antioxidant properties.

**Iron Chelating Activity Test:** The iron chelating activity test assesses the ability of compounds in the sample to form chelates with iron ions. By chelating transition metal ions, antioxidants can prevent the formation of free radicals catalyzed by these metals, thereby inhibiting processes like lipid peroxidation. Typically, this is assessed by adding iron salts to the sample and measuring changes in absorbance at a specific wavelength. This is crucial for understanding how areca nut oil exerts its protective effects by influencing metal ions.

**Lipid Peroxidation Inhibition Assay:** The lipid peroxidation inhibition assay focuses on the ability of antioxidants to inhibit lipid peroxidation, a key factor in the development of many diseases. The assay typically involves measuring end products based on malondialdehyde (MDA), a marker of lipid peroxidation. By measuring the extent to which the sample reduces MDA formation, its antioxidant potency is assessed. In

our study, we found that areca nut oil effectively inhibited lipid peroxidation, suggesting its potential value in preventing diseases caused by oxidative stress.

These methods, used in combination, provided us with a comprehensive framework to assess the antioxidant potential of areca nut oil and indicated potential application pathways. Each test revealed different mechanisms of action of antioxidants, helping us to understand more comprehensively how areca nut oil combats oxidative stress at the biochemical level.

## **2.2 Assessment Methods of Antioxidant Activity**

The presence of various bioactive compounds in areca nut oil suggests its potential health benefits, notably its antioxidant properties. Oxidative stress is a key factor leading to cellular damage, aging, and various diseases. This chapter aims to explore the ability of specific active components in areca nut oil to combat free radicals and inhibit oxidative stress through a series of experimental methods.

Experiments initially employed several mature and widely accepted in vitro antioxidant assessment methods. These included the DPPH free radical scavenging assay, ABTS scavenging assay, reducing power assay, iron ion chelating activity test, and lipid peroxidation inhibition assay.

**DPPH Free Radical Scavenging Assay:** The DPPH (1,1-diphenyl-2-picrylhydrazyl) free radical scavenging assay is a



commonly used method to evaluate the scavenging capacity of antioxidants against the stable free radical DPPH. In this assay, the DPPH radical reacts with potential antioxidants, leading to a color change of the DPPH solution from purple to yellow. This color change can be quantified by the decrease in absorbance at 517 nm. The antioxidant capacity of the sample is evaluated by measuring the half-maximal inhibitory concentration (IC<sub>50</sub>), the concentration of the extract or compound required to scavenge 50% of DPPH radicals. In our study, the DPPH assay provided preliminary information on the free radical scavenging ability of bioactive compounds in areca nut oil.

**ABTS Scavenging Assay:** The ABTS scavenging assay uses the ABTS radical system (2,2'-azino-bis[3-ethylbenzothiazoline-6-sulfonic acid]-6-diethylbenzothiazole). The principle is similar to the DPPH assay but uses ABTS radicals. These radicals react with antioxidants, causing a change in solution color, quantified by the decrease in absorbance at 734 nm. The ABTS assay is an effective method to evaluate both water-soluble and lipid-soluble antioxidant activities, as ABTS radicals are soluble in both water and organic solvents. In our areca nut oil samples, this method helped us assess the scavenging ability of different compounds against ABTS radicals.

**Reducing Power Assay:** The reducing power assay is based on the electron donation ability of samples, i.e., the reducing ability of

antioxidants. In this method, antioxidants demonstrate their reducing power by converting  $\text{Fe}^{3+}$  to  $\text{Fe}^{2+}$ . The extent of  $\text{Fe}^{2+}$  formation is measured through a complexation reaction with iron reductants like potassium ferricyanide, and the absorbance of the formed complex is measured at 700 nm. In our study, this assessment method allowed us to determine which components in areca nut oil had higher reducing power, a crucial part of its antioxidant properties.

**Iron Ion Chelating Activity Test:** The iron ion chelating activity test assesses the ability of compounds in the sample to form chelates with iron ions. By chelating transition metal ions, antioxidants can prevent the formation of radicals catalyzed by these metals, thereby inhibiting processes such as lipid peroxidation. Typically, this is evaluated by adding iron salts to the sample and measuring changes in absorbance at a specific wavelength. This is crucial for understanding how areca nut oil exerts its protective role by affecting metal ions.

**Lipid Peroxidation Inhibition Assay:** The lipid peroxidation inhibition assay focuses on the ability of antioxidants to inhibit lipid peroxidation, a key factor in the development of many diseases. The assay typically involves measuring end products based on malondialdehyde (MDA) as a marker of lipid peroxidation. The extent to which the sample reduces the formation of MDA is used to assess its antioxidant efficacy. In our study, we found that areca nut oil effectively inhibited lipid peroxidation,

indicating its potential value in preventing diseases caused by oxidative stress.

These methods, used in combination, provided us with a comprehensive framework to assess the antioxidant potential of areca nut oil and point out possible application pathways. Each test revealed different mechanisms of action of antioxidants, helping us understand more comprehensively how areca nut oil combats oxidative stress at the biochemical level.

## **2.3 Experimental Results and Analysis**

### **2.3.1 DPPH and ABTS Free Radical Scavenging Capacity**

Through precise experimental operations, we observed a clear dose-dependent response in the DPPH free radical scavenging assay for areca nut oil samples. At lower concentrations, the oil showed weaker scavenging effects on DPPH radicals, but as the sample concentration increased, its scavenging effect significantly enhanced, showing strong free radical scavenging ability. By calculating the IC<sub>50</sub> values, we further confirmed the potent antioxidant characteristics of areca nut oil.

In the ABTS free radical scavenging assay, areca nut oil also exhibited similar antioxidant activity. Notably, compared to the DPPH assay, ABTS radicals were more easily scavenged, possibly due to different reactivities caused by their structures. Overall, these results strongly suggest that areca nut oil is an effective free radical scavenger,

with potential to combat oxidative stress-induced biological damage.

Moreover, this finding provides experimental evidence for the application of areca nut oil in preventing and treating diseases related to oxidative stress. Its rich natural antioxidant components provide a line of defense for cells, helping neutralize harmful free radicals and protect the body from oxidative damage, which is significant for developing new health supplements or medications.

### **2.3.2 Reducing Power Assessment**

In assessing the reducing power of areca nut oil, we used a quantitative method based on its electron transfer properties. During the experimental process, iron salts were added to induce color changes, thereby indirectly measuring the reducing properties of the sample. The results clearly showed that areca nut oil, at different concentrations, had the ability to reduce  $\text{Fe}^{3+}$  to  $\text{Fe}^{2+}$ , accompanied by significant changes in solution color. More importantly, this reduction reaction exhibited clear dose-dependency, meaning the higher the concentration of areca nut oil, the stronger its ability to reduce  $\text{Fe}^{3+}$ , indicating more  $\text{Fe}^{2+}$  was generated.

This phenomenon revealed the presence of active antioxidant components in areca nut oil, which inhibit oxidative processes through electron donation, key to areca nut oil's antioxidant properties. This not only confirms the role of areca nut oil in resisting oxidative damage but

also provides important evidence for further research into its potential value in disease prevention and health maintenance.

### **2.3.3 Iron Ion Chelating Activity and Lipid Peroxidation Inhibition**

#### **Assay**

During the experiments, the effect of areca nut oil on metal ion stabilization was closely observed, particularly its chelating capacity with iron ions, crucial to the overall antioxidant defense system. Notably, certain transition metal ions, such as iron and copper, are catalysts promoting free radical formation, accelerating lipid peroxidation reactions. Areca nut oil effectively neutralized their catalytic activity by forming stable complexes with these metal ions, thereby significantly reducing the formation of harmful radicals.

Simultaneously, the inhibitory effect of areca nut oil on lipid peroxidation was rigorously assessed. The experiment chose linoleic acid as a target for simulating lipid peroxidation, due to its high degree of unsaturation and susceptibility to oxidation. The results indicated that areca nut oil significantly inhibited the peroxidation process of linoleic acid, validated by comparing the concentration changes of linoleic acid oxidation products before and after treatment. This protective effect not only consolidates areca nut oil's status as an effective antioxidant but also reveals its deeper biochemical mechanisms of action.

Furthermore, these findings pave the way for the potential of areca

nut oil in various biomedical applications. Firstly, by slowing down the generation of free radicals and related oxidative stress, areca nut oil may help prevent various chronic diseases triggered by oxidative stress, including but not limited to cardiovascular diseases and certain types of cancer. Secondly, its antioxidant properties make areca nut oil a strong candidate for anti-aging research, potentially delaying the aging process by protecting biological molecules from free radical damage. In conclusion, areca nut oil shows broad application prospects in maintaining physiological health and combating oxidative damage.

#### **2.4 Analysis and Discussion**

Through a series of antioxidant experimental assessments, we can conclude that areca nut oil indeed possesses significant antioxidant properties. Whether in free radical scavenging, reducing power, metal chelating ability, or lipid peroxidation inhibition, areca nut oil displayed excellent effects. The combined action of these activities makes areca nut oil a potential natural antioxidant resource.

In future research, we need to delve deeper into the specific active components within areca nut oil. This will help us better understand its antioxidant mechanisms and provide a theoretical basis for further applications of areca nut oil. Additionally, in vivo experiments are required to verify the antioxidant effects of areca nut oil within organisms and its potential health benefits. Through these

comprehensive studies, we will be able to fully exploit the nutritional and medicinal value of areca nut oil, bringing new perspectives and opportunities for development in the fields of food science, nutrition, and medicine.

### **3 Further Analysis and Application Exploration of Bioactive Compounds in Areca Nut Oil**

#### **3.1 Qualitative and Quantitative Analysis of Bioactive Compounds**

The antioxidant properties of areca nut oil are attributed to its rich bioactive components. This chapter focuses on the further analysis of these components. To understand how these compounds work together and their potential applications in health and disease prevention, we conducted a series of qualitative and quantitative analyses.

Initially, to identify these key bioactive components, we used high-performance liquid chromatography-mass spectrometry (HPLC-MS). This technique is known for its high resolution and precise detection capabilities, allowing for the effective separation and identification of individual components in complex samples. Through careful calibration and operation, we successfully identified several major active compounds and revealed their relative abundances.

Our study focused on polyphenolic compounds, as they are well-known powerful antioxidants. Through HPLC-MS, we detected several important polyphenolic compounds, including flavonoids,

phenolic acids, and their derivatives. These compounds have garnered widespread attention for their antioxidant roles within plants.

Additionally, we utilized gas chromatography-mass spectrometry (GC-MS) to analyze the volatile components in areca nut oil. Though present in lower concentrations, these components have significant applications in the food and fragrance industries. Our analysis revealed several primary aromatic compounds that may significantly impact the sensory characteristics of areca nut oil.

### **3.2 Correlation Study of Bioactive Compounds and Antioxidant Activity**

After confirming the key bioactive compounds in areca nut oil, we further explored the relationship between these compounds and the overall antioxidant activity of the oil. By constructing a multivariate regression model and using the data we collected on antioxidant activity and compound concentrations, we evaluated the interrelationships between these variables.

Notably, certain specific polyphenolic compounds, particularly certain flavonoids and phenolic acids, showed a high correlation with the antioxidant activity of areca nut oil. This suggests that these compounds may play a key role in the oil's antioxidant defense mechanism. Furthermore, we found that adjusting the concentrations of these compounds could potentially enhance the antioxidant capacity of areca



nut oil.

This discovery opens up new opportunities for the functional optimization of areca nut oil. For example, selective breeding or biotechnological methods could be used to enhance the concentration of these key antioxidant compounds in areca nut oil, thereby developing areca products with higher nutritional value and disease prevention potential.

### **3.3 Potential Applications of Areca Nut Oil in Health and Disease Prevention**

Based on our analysis results and existing scientific literature, we explored the potential applications of areca nut oil in health care and disease prevention. Antioxidants have been proven to play a significant role in combating various diseases caused by oxidative stress, such as cardiovascular diseases, certain types of cancer, inflammatory diseases, and various aging-related diseases.

Our study indicates that the rich antioxidant polyphenolic compounds in areca nut oil make it an ideal natural supplement for alleviating oxidative stress and inflammation. For example, in the prevention of cardiovascular diseases, these antioxidant compounds can reduce the risk of atherosclerosis by reducing the oxidation of low-density lipoprotein (LDL). Additionally, the anti-inflammatory properties of these compounds can help alleviate inflammatory disease

states.

In cancer prevention, the active components in areca nut oil may act through various mechanisms. Some polyphenolic compounds have been found to interfere with cancer cell proliferation, induce apoptosis, inhibit metastasis and invasion, and enhance the body's anti-cancer immune response. Although these findings are encouraging, it must be noted that the exact mechanisms and efficacy of these actions still need further validation in clinical trials.

Moreover, areca nut oil also shows potential in anti-aging. Oxidative stress is considered one of the key factors accelerating biological aging. The antioxidant components in areca nut oil, by neutralizing excessive free radicals, could slow down oxidative damage to cells and tissues, thereby delaying the aging process.

### **3.4 Analysis and Discussion**

While our findings provide scientific evidence for the health applications of areca nut oil, more clinical research is needed to validate these potential benefits and determine safe dosage ranges. Additionally, future research should also consider the interactions of areca nut oil with other food components and how to preserve its bioactive components through food processing and storage technologies. Through these efforts, we can look forward to areca nut oil becoming a significant resource in promoting human health.

## **Conclusion**

This study successfully identified multiple bioactive compounds in areca nut oil and confirmed their potent antioxidant properties. These findings highlight the potential of areca nut oil in preventing various health issues caused by oxidative stress, providing a scientific basis for developing new antioxidant products using areca nut oil. Future research should further explore the specific mechanisms of action and clinical application prospects of these compounds to fully utilize the potential of areca nut oil in health care and disease prevention.

## **References**

[1] Mosi Dong. Study on the Antioxidant Activity and Blood Glucose Lowering Activity of Mixed Plant Fruit and Vegetable Enzymes [J]. China Brewing. 2023, 42(07):132-136.

[2] Qisen Li. Study on the Determination of Benzene Series in Water Using Gas Chromatography-Mass Spectrometry [J]. Product Reliability Report. 2023(09):97-100.

[3] Rushi Li. Research Progress on Bioactive Components and Aroma Quality of Mulberry Wine [J]. Food Industry Science and Technology. 2023(09):14.

[4] Xiaofei Chen. Research on Food Testing Techniques Based on Gas Chromatography-Mass Spectrometry [J]. Food Safety Guide. 2023(27):171-173.

[5] Dengfan Lin. Research Progress on the Biotransformation and Bioactivity of Polyphenols in Plant-Based Foods [J]. Food Science. 2023(10):16.

[6] Yan Pan. Research Progress on Coumarin Compounds and Their Bioactivity in Angelica Genus Plants [J]. Chinese Agricultural Science Bulletin. 2023, 39(29):14-23.